

***Ranunculus maoxianensis* (Ranunculaceae), a new species from northwestern Sichuan, China, with an emended description of *R. chongzhouensis*, the putative closest ally of the new species**

Wen-Qun Fei^{1,2}, Qiong Yuan^{1,3}, Qin-Er Yang^{1,3}

1 Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510655, Guangdong, China **2** University of Chinese Academy of Sciences, Beijing 100049, China **3** Center of Conservation Biology, Core Botanical Gardens, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510655, Guangdong, China

Corresponding author: Qiong Yuan (yuanqiong@scib.ac.cn)

Academic editor: M. Pellegrini | Received 19 October 2022 | Accepted 6 January 2023 | Published 1 February 2023

Citation: Fei W-Q, Yuan Q, Yang Q-E (2023) *Ranunculus maoxianensis* (Ranunculaceae), a new species from northwestern Sichuan, China, with an emended description of *R. chongzhouensis*, the putative closest ally of the new species. *PhytoKeys* 219: 77–96. <https://doi.org/10.3897/phytokeys.219.96510>

Abstract

Ranunculus maoxianensis (Ranunculaceae), a new species from Jiuding Shan in Maoxian county, northwestern Sichuan province, China, is here illustrated and described. The species is morphologically similar to *R. chongzhouensis*, a species also occurring in Sichuan, in having reniform leaves and puberulous receptacles, carpels and achenes, but differs by having leaves adaxially puberulous with shorter appressed hairs 0.16–0.28 mm long (vs. longer appressed hairs 0.55–0.85 mm long), larger flowers (1.8–2 cm vs. 1.4–1.6 cm in diameter), larger (8–10 × 5.5–6.5 mm vs. 6–7 × 4.5–5 mm) and widely obovate petals (vs. obovate), more numerous stamens (35–55 vs. 12–18), and subglobose gynoeceum and aggregate fruit (vs. ellipsoid). The two species are also different in chromosome number and chromosome morphology. *Ranunculus chongzhouensis* has a karyotype of $2n = 2x = 16 = 10m + 6sm$ while *R. maoxianensis* has a karyotype of $2n = 4x = 32 = 16m + 16sm$. An emended description of *R. chongzhouensis* is provided, and its geographical distribution is largely extended.

Keywords

Asia, buttercups, chromosome number, *Ranunculus* sect. *Ranunculus*

Introduction

Ranunculus L., comprising approximately 600 species, is the largest genus in the Ranunculaceae and is widely distributed in all continents (Tamura 1995; Hörandl et al. 2005; Paun et al. 2005; Hörandl and Emadzade 2012). In China, one of the centers of species diversity in *Ranunculus*, more than 150 species and 30 varieties are currently recognized in the genus (Wang 1995a, b, 1996, 2007, 2008, 2013, 2015, 2016, 2018, 2019a, b, 2022; Yang 2000; Wang and Gilbert 2001; Wang and Liao 2009; Luo and Zhao 2013; Wang and Chen 2015; Wang et al. 2016; Yuan and Yang 2017a, b, c; Zhang et al. 2020; Fei et al. 2022, 2023a, b). Many taxa in the genus occur in the Hengduan Mountains region in southwestern China, including mainly southeastern Gansu, eastern Qinghai, western Sichuan, southeastern Xizang (Tibet) and northwestern Yunnan, one of the most important biodiversity hotspots worldwide (Zhang et al. 2009; Sun et al. 2017). In total, 56 species and nine varieties of *Ranunculus* are known from that region (Wang 1993, 1995a, b, 2008, 2013, 2015, 2022; Yang 2000; Wang and Gilbert 2001; Wang and Liao 2009; Wang and Chen 2015; Yuan and Yang 2017c).

During a botanical expedition to the Hengduan Mountains region from June to August 2022 for the first author's Ph.D. dissertation project, we discovered an unusual population (Figs 1, 2) of *Ranunculus* on Jiuding Shan in Maoxian county, northwestern Sichuan province, China. The plants are somewhat similar to *R. chongzhouensis* W.T. Wang (Figs 3–10), a species also occurring in Sichuan, in having reniform leaves and puberulous receptacles, carpels and achenes, but differ by an array of characters, such as the length of hairs on the adaxial side of leaf blades, size of flowers, size and shape of petals, number of stamens, and shape of the gynoeceum and aggregate fruit. Moreover, our chromosomal examination revealed that the Maoxian population in question has a chromosome number of $2n = 16$ (Fig. 11A), while *R. chongzhouensis* has a chromosome number of $2n = 32$ (Fig. 11B). Therefore, we determined that this population represents a hitherto undescribed species, which we describe as *R. maoxianensis* below. Furthermore, we found that *R. chongzhouensis*, the putative closest ally of *R. maoxianensis*, is much more widely distributed than documented before and that its original description is not complete, lacking a description of root and floral characters, with the description of indumentum of leaf blades, receptacles and achenes being also incorrect. Based on our critical observations of herbarium specimens and living plants in the wild, the description of this species is here emended. We also largely extended its geographical distribution.

Materials and methods

For morphological comparison, we critically examined specimens or high-resolution specimen images of *Ranunculus* at CDBI, KUN, PE, and WCSBG (acronyms according to Thiers 2022). We also observed living plants in three populations of *R. chongzhouensis* from Sichuan and one population of the new species *R. maoxianensis* (Table 1) at flowering and fruiting time (June to July). We observed characters of roots, stems, leaves, pedicels, flowers, receptacles, petals, stamens, gynoeceum, carpels,

Table 1. Information about three populations of *Ranunculus chongzhouensis* and one of *R. maoxianensis* sp. nov. observed in the wild. Populations with an asterisk were used for SEM observation of the leaf epidermis and chromosomal examination.

Taxon	Voucher	Locality
<i>R. chongzhouensis</i>	W.Q. Fei 915 (IBSC)	China, Sichuan, Chongzhou, Jiguan Shan
	W.Q. Fei 577 (IBSC)*	China, Sichuan, Dayi, Xiling Xue Shan
	W.Q. Fei & H.S. Wu 395 (IBSC)	China, Sichuan, Xiaojin, Siguniang Shan
<i>R. maoxianensis</i>	W.Q. Fei 565 (IBSC)*	China, Sichuan, Maoxian, Jiuding Shan

aggregate fruit and achenes, paying special attention to the indumentum of basal leaves, size of flowers, size and shape of petals, number of stamens, and the shape of gynoecium and aggregate fruit.

For scanning electron microscopy (SEM), dry leaves were taken from herbarium specimens (Table 1) and mounted directly onto stubs using double-sided sellotape, gold-coated, and then observed and photographed under a JSM-6360LV scanning electron microscope.

For chromosomal examination, living plants of *Ranunculus chongzhouensis* from Xiling Xue Shan in Dayi, Sichuan, and *R. maoxianensis* from its type locality, i.e., Jiuding Shan in Maoxian, Sichuan (Table 1), were cultivated in pots in the experimental garden of South China Botanical Garden, Chinese Academy of Sciences. Root tips were pretreated in 0.1% colchicine for 2.5 h, fixed in Carnoy I (glacial acetic acid: absolute ethanol = 1: 3), then macerated in 1 M HCl at 37 °C for 45 min, and stained and squashed in Carbol fuchsin. Karyotype formulas were based on the data of measurements of mitotic-metaphase chromosomes of three cells taken from photographs. We followed the acronyms proposed by Levan et al. (1964) to describe the karyotypes.

Results and discussion

Our critical observations on herbarium specimens and living plants in the wild indicate that *Ranunculus maoxianensis* (Figs 1, 2) is morphologically similar to *R. chongzhouensis* (Figs 3–10) in having reniform leaves and puberulous receptacles, carpels and achenes, but differs by having leaves adaxially puberulous with shorter appressed hairs 0.16–0.28 mm long (vs. longer appressed hairs 0.55–0.85 mm long) (Fig. 12A, C), larger flowers (1.8–2.0 cm vs. 1.4–1.6 cm in diameter), larger (8–10 × 5.5–6.5 mm vs. 6–7 × 4.5–5 mm) and widely obovate petals (vs. obovate), more numerous stamens (35–55 vs. 12–18), and subglobose gynoecium and aggregate fruit (vs. ellipsoid). The difference in the indumentum of leaves between the two species is further confirmed by our SEM results (Fig. 12B, D). A detailed morphological comparison between *R. maoxianensis* and *R. chongzhouensis* is given in Table 2.

Our chromosomal examination reveals that *Ranunculus maoxianensis* is a diploid species with $2n = 2x = 16 = 10m + 6sm$ (Fig. 11A, C), while *R. chongzhouensis* is a tetraploid with $2n = 4x = 32 = 16m + 16sm$ (Fig. 11B, D). This result lends strong support to the description of *R. maoxianensis* as a new species.

Table 2. Morphological comparison between *Ranunculus chongzhouensis* and *R. maoxianensis* sp. nov.

	<i>R. chongzhouensis</i>	<i>R. maoxianensis</i>
Stems	10–25 cm tall	25–55 cm tall
Basal leaves	5–8, blades 2.2–3.1 × 3.1–3.7 cm, chartaceous, adaxially appressed puberulous with hairs 0.55–0.85 mm long, abaxially glabrous or sometimes appressed puberulous	2–5, blades 2.2–3.2 × 3.8–5.2 cm, herbaceous, adaxially appressed puberulous with hairs 0.16–0.28 mm long, abaxially appressed puberulous
Flowers	terminal, 4–10, 1.4–1.6 cm in diameter	terminal, 4–10, 1.8–2 cm in diameter
Receptacles	3–5 mm long, clavate, puberulous	3.5–4 mm long, clavate, puberulous
Petals	6–7 × 4.5–5 mm, obovate	8–10 × 5.5–6.5 mm, widely obovate
Stamens	12–18	35–55
Gynoecium	ellipsoid	subglobose
Carpels	20–40; ovaries ovoid or widely ovoid puberulous, styles ca. 0.9 mm long, glabrous, slightly recurved at apex	16–22; ovaries ovoid or widely ovoid, puberulous, styles ca. 0.9 mm long, glabrous, straight or apex recurved
Aggregate fruit	ellipsoid	subglobose
Achenes	ca. 2 × 1.5 mm, obliquely or widely ovoid, puberulous, styles ca. 1 mm long, straight or apex recurved.	ca. 2.5 × 2 mm, obliquely or widely ovoid, puberulous, styles ca. 1 mm long, straight or apex recurved.

Our literature consultation and critical observations on herbarium specimens and living plants in the wild reveal that *Ranunculus chongzhouensis*, the putative closest ally of *R. maoxianensis*, lacks the description of root and floral characters, with the description of indumentum of leaf blades, receptacles and achenes by Wang (2015) being also incorrect. Wang (2015) described this species based on a single specimen, *Z.B. Feng, D.H. Zhu & X.J. Li 4171* (PE; Fig. 3A), from Jiguan Shan in the Anzihe Nature Reserve in Chongzhou city, Sichuan province, China. The three plants on the sheet are all fruiting, lacking roots and flowers. Therefore, the number of basal leaves and floral morphology were not mentioned in the original description of *R. chongzhouensis*. We traced two isotype sheets of *R. chongzhouensis* from WCSBG (Fig. 3B, C), which Wang (2015) did not see when he described this species as new. Wang (2015) described the basal leaves of *R. chongzhouensis* as adaxially glabrous and abaxially appressed puberulous, and the receptacles and achenes as glabrous. However, the basal leaves of this species are adaxially appressed puberulous (Fig. 3D, E) and abaxially glabrous (Fig. 3F, G), and the receptacles (Fig. 3H) and achenes (Fig. 3H, I) are puberulous. These results are further confirmed by our observations of living plants in three populations of *R. chongzhouensis*, respectively, from Chongzhou (type locality), Dayi, and Xiaojin, all in Sichuan province (Figs 4–9). In addition, we found that the indumentum on the abaxial side of the leaf blades of *R. chongzhouensis* is somewhat variable between populations. The leaf blades are often abaxially glabrous (Figs 4E, 6E), but sometimes abaxially puberulous (Fig. 8E).

From our survey of herbarium specimens and fieldwork, we found that *Ranunculus chongzhouensis* is much more widely distributed than reported by Wang (2015). In addition to its type locality, this species is also distributed in Baoxing, Dayi, Heishui, Luding, Songpan, and Xiaojin counties. Selected specimens from Baoxing (Fig. 10A), Heishui (Fig. 10B), Luding (Fig. 10C) and Songpan (Fig. 10D) are shown in Fig. 10.

Ranunculus maoxianensis is readily assigned to *R. sect. Ranunculus* due to its swollen achenes with a distinct beak and receptacles hardly enlarged after anthesis. Wang (2015) also included *R. chongzhouensis* in this section. We accept the sectional placement of this species.

Taxonomic treatment

Ranunculus maoxianensis W.Q.Fei, Q.Yuan & Q.E.Yang, sp. nov.

urn:lsid:ipni.org:names:77313248-1

Figs 1, 2

Diagnosis. *Ranunculus maoxianensis* is similar to *R. chongzhouensis* in leaf blade shape and indumentum of the receptacles, carpels and achenes. However, it is easily distinguishable by having leaves adaxially puberulous with shorter appressed hairs 0.16–0.28 mm long (vs. longer appressed hairs 0.55–0.85 mm long), larger flowers (1.8–2 cm vs. 1.4–1.6 cm in diameter), larger (8–10 × 5.5–6.5 mm vs. 6–7 × 4.5–5 mm) and widely obovate petals (vs. obovate), more numerous stamens (35–55 vs. 12–18), and subglobose gynoecium and aggregate fruit (vs. ellipsoid).

Type. CHINA. Sichuan: Maoxian, Nanxin town, Jiuding Shan, 31°30'36.28"N, 103°46'52.01"E, alt. 3274 m, in *Rhododendron* forests, 7 June 2022, W.Q. Fei 565 (holotype: IBSC; isotypes: CDBI, IBSC, PE).

Description. *Herb* perennial, terrestrial. *Roots* fibrous, slender. *Stems* 25–55 cm tall when in bloom, branched, erect, sparsely puberulous. *Basal leaves* 2–5, long petiolate; petioles 7–20 cm long, sparsely puberulous; blades 2.2–3.2 × 3.8–5.2 cm, reniform, 3-lobed or 3-partite, herbaceous, adaxially green, appressed puberulous with hairs 0.16–0.28 mm long, abaxially light green, puberulous with hairs 0.65–1.1 mm long, base truncate or cordate, central segment 1.2–1.5 × 0.8–1.2 cm, rhombic or rhombic-obovate, margin crenulate, lateral segments 1.5–1.8 × 2–2.6 cm, obliquely flabellate, unequally 2-partite, margin crenulate. *Lower cauline leaves* 1–2, similar to basal ones but smaller. *Upper cauline leaves* 2–4, subsessile or sessile, 3-sected, segments 1.5–3 × 0.3–0.8 mm, obtriangular, lanceolate or linear, entire or 3–5-lobed. *Inflorescences* terminal, 4–10-flowered. *Flowers* 1.8–2 cm in diameter; pedicels 5–10 cm long, appressed puberulous; receptacles 3.5–4 mm long, clavate, puberulous; sepals 5, 4.6–5 × 2–3 mm, elliptic to obovate, green tinged with yellowish, adaxially glabrous, abaxially puberulous; petals 5(–6), 8–10 × 5.5–6.5 mm, widely obovate, yellow, glabrous, apex rounded or subtruncate, nectary pit without a scale, claw ca. 0.6 mm long; stamens 35–55, filaments 1.5–2 mm long, narrowly linear, anthers 1–1.2 mm long, oblong; gynoecium subglobose; carpels 16–22, ovaries ca. 0.9 × 0.8 mm, ovoid or widely ovoid, laterally flattened, biconvex, puberulous, styles ca. 0.9 mm long, glabrous, straight or apex recurved. *Aggregate fruit* ca. 7 × 7 mm, subglobose; achenes ca. 2.5 × 2 mm, obliquely or widely ovoid, laterally flattened, biconvex, puberulous, styles ca. 1 mm long, persistent, glabrous, straight or apex recurved.

Etymology. The specific epithet refers to the type locality of the new species, i.e. Maoxian county in northwestern Sichuan province, China.

Phenology. Flowering from June to July; fruiting from July to August.

Distribution and habitat. *Ranunculus maoxianensis* is currently known only from its type locality, i.e., Jiuding Shan in Maoxian county, northwestern Sichuan province, China (Fig. 13). It grows in *Rhododendron* forests at altitudes of 3200–3400 m above sea level.

Conservation status. Although *Ranunculus maoxianensis* is currently known only from its type locality, i.e., Jiuding Shan in Maoxian county, northwestern Sichuan

province, China, our observations on living plants at its type locality revealed that this species is very common in *Rhododendron* forests at altitudes of 3200–3400 m above sea level. It should better be categorized as “Least Concern (LC)”, following the IUCN Standards and Petitions Committee (IUCN 2022).

Additional specimens examined (paratype). CHINA. Sichuan: Maoxian, W.Q. Fei & H.S. Wu 397 (IBSC).

***Ranunculus chongzhouensis* W.T. Wang, Bull. Bot. Res., Harbin 35(5): 645. 2015**
Figs 3–10

Type. CHINA. Sichuan: Chongzhou city, Anzihe Nature Reserve, Jiguan Shan, alt. 3000 m, in forests, 27 July 2007, Z.B. Feng, D.H. Zhu & X.J. Li 4171 (holotype: PE!; isotypes: WCSBG!).

Description. *Herbs* perennial, terrestrial. *Roots* fibrous, slender. *Stems* 10–25 cm tall, ascending or erect, branched, puberulous. Basal leaves 5–8, 3-lobed or 3-partite, long petiolate; petioles 4–10 cm long, sparsely puberulous; blades 2.2–3.1 × 3.1–3.7 cm, reniform, chartaceous, adaxially green, appressed puberulous with hairs 0.55–0.85 mm long, abaxially light green, glabrous, sometimes puberulous, base cordate, central segment 0.6–1 × 0.9–1.4 cm, obtrapezoid or obovate-obtrapezoid, margin 3-crenulate, lateral segments 0.8–1.2 × 1.5–2.3 cm, obliquely flabellate, unequally 2-lobed, margin crenulate. *Lower cauline leaves* 1 or absent, similar to basal ones but smaller. *Upper cauline leaves* 2–3, 3-sected, subsessile, segments 1.1–1.5 × 0.3–0.9 mm, obliquely flabellate, lanceolate to linear, margin entire or 3–4-lobed. *Inflorescences* terminal, 4–10-flowered. *Flowers* 1.4–1.6 cm in diameter; pedicels 5–10 cm long, appressed puberulous; receptacles 3–5 mm long, clavate, puberulous; sepals 5, 3.9–4.5 × 2.5–3 mm, elliptic to obovate, green tinged with yellowish, adaxially glabrous, abaxially puberulous; petals 5(–6), 6–7 × 4.5–5 mm, obovate, yellow, glabrous, apex truncate or subtruncate, nectary pit without a scale, claw ca. 0.5 mm long; stamens 12–18, filaments ca. 2 mm long, narrowly linear, anthers ca. 1 mm long, oblong; gynoecium ellipsoid; carpels 20–40, ovaries ca. 0.9 × 0.8 mm, ovoid or widely ovoid, laterally flattened, biconvex, puberulous, styles ca. 0.9 mm long, glabrous, apex slightly recurved. *Aggregate fruit* ca. 7 × 5 mm, ellipsoid; achenes ca. 2 × 1.5 mm, obliquely or widely ovoid, laterally flattened, biconvex, puberulous, styles ca. 1 mm long, persistent, straight or apex recurved.

Phenology. Flowering from June to July; fruiting from July to August.

Distribution and habitat. *Ranunculus chongzhouensis* is distributed in Baoxing, Chongzhou, Dayi, Heishui, Luding, Songpan, and Xiaojin in Sichuan province, China (Fig. 13). It grows in forests or meadows at elevations of 2900–4150 m above sea level.

Additional specimens examined. CHINA. Sichuan: Baoxing, W.Q. Fei & H.S. Wu 371 (IBSC); Chongzhou, W.Q. Fei 915 (IBSC), W.B. Ju, L. Zhang & D.K. Chen AZH01296 (CDBI); Dayi, W.Q. Fei 577 (IBSC), J.P. Luo & H.M. Li 613 (IBSC), Y.P. Zeng, Y.F. Luo & Y.Q. Tao 149 (IBSC); Heishui, W.Q. Fei 719 (IBSC); Luding, W.Q. Fei 754 (IBSC); Songpan, W.Q. Fei 725 (IBSC); Xiaojin, W.Q. Fei & H.S. Wu 395 (IBSC).

Figures




Figure 1. *Ranunculus maoxianensis* sp. nov. in the wild (China, Sichuan, Maoxian) **A** habitat **B** habit **C** roots **D** portion of stem **E** leaf blade (left: adaxial side; right: abaxial side) **F** flower (lateral view) **G** flower (top view) **H** sepal (left: abaxial side; right: adaxial side) **I** petal (left: adaxial side; right: abaxial side) **J** stamens **K** carpels **L** aggregate fruit **M** achenes **N** receptacle. Photographed by Wen-Qun Fei.



Figure 2. Holotype (A) and isotype (B–D) sheets of *Ranunculus maoxianensis* sp. nov.




Figure 3. Holotype (A) and isotype (B, C) sheets of *Ranunculus chongzhouensis*, showing the general morphology and indumentum of leaf blade (D–G), aggregate fruit (H, I) and receptacle (H). D, E portion of adaxial side of leaf blade (appressed puberulous with longer hairs) from A and B respectively; E, G portion of abaxial side of leaf blade (glabrous) from A and B respectively; H, I portion of aggregate fruit (puberulous) from A and C respectively; H portion of the receptacle (puberulous) from A.




Figure 4. *Ranunculus chongzhouensis* in the wild (China, Sichuan, Chongzhou, the type locality) **A** habitat **B** habit **C** roots **D** portion of stem **E** leaf blade (left: adaxial side; right: abaxial side) **F** flower (lateral view) **G** flower (top view) **H** sepal (left: abaxial side; right: adaxial side) **I** petal (left: adaxial side; right: abaxial side) **J** stamens **K** carpels **L** aggregate fruit **M** achenes **N** receptacle. Photographed by Wen-Qun Fei.



Figure 5. Selected specimens of *Ranunculus chongzhouensis* from its type locality, i.e., Chongzhou city in Sichuan province, China **A, B** W.Q. Fei 915 (IBSC).




Figure 6. *Ranunculus chongzhouensis* in the wild (China, Sichuan, Dayi) **A** habitat **B** habit **C** roots **D** portion of stem **E** leaf blade (left: adaxial side; right: abaxial side) **F** flower (lateral view) **G** flower (top view) **H** sepal (left: abaxial side; right: adaxial side) **I** petal (left: adaxial side; right: abaxial side) **J** stamens **K** carpels **L** aggregate fruit **M** achenes **N** receptacle. Photographed by Wen-Qun Fei.



Figure 7. Selected specimens of *Ranunculus chongzhouensis* from Dayi county in Sichuan province, China **A, B** W.Q. Fei 577 (IBSC).




Figure 8. *Ranunculus chongzhouensis* in the wild (China, Sichuan, Xiaojin) **A** habitat **B** habit **C** roots **D** portion of stem **E** leaf blade (left: adaxial side; right: abaxial side) **F** flower (lateral view) **G** flower (top view) **H** sepal (left: abaxial side; right: adaxial side) **I** petal (left: adaxial side; right: abaxial side) **J** stamens **K** carpels **L** aggregate fruit **M** achenes **N** receptacle. Photographed by Wen-Qun Fei.



Figure 9. Selected specimens of *Ranunculus chongzhouensis* from Xiaojin county in Sichuan province, China **A, B** W.Q. Fei 395 (IBSC).




Figure 10. Selected specimens of *Ranunculus chongzhouensis* from Baoxing (A), Heishui (B), Luding (C) and Songpan (D) in Sichuan province, China **A** W.Q. Fei 371 (IBSC) **B** W.Q. Fei 719 (IBSC) **C** W.Q. Fei 754 (IBSC) **D** W.Q. Fei 725 (IBSC).




Figure 11. Mitotic metaphase chromosomes (**A, B**) and karyotypes (**C, D**) of *Ranunculus maoxianensis* sp. nov. (**A, C**) and *R. chongzhouensis* (**B, D**), with m = median-centromeric chromosome, sm = submedian-centromeric chromosome.




Figure 12. Portion of the adaxial side of the leaf blade of *Ranunculus chongzhouensis* (**A, B**) and *R. maoxianensis* sp. nov. (**C, D**), showing the difference in length of hairs. The hairs on the adaxial side of the leaf blade in *R. chongzhouensis* are 0.55–0.85 mm long, and those in *R. maoxianensis* are 0.16–0.28 mm long **A, C** photographed in the wild and **B, D** photographed with SEM.




Figure 13. Distribution of *Ranunculus chongzhouensis* (■) and *R. maoxianensis* (●). Black arrow indicates the type locality of *R. chongzhouensis*, i.e., Chongzhou city in Sichuan province, China.

Acknowledgements

We are grateful to one anonymous reviewer, Dr. Marco Pellegrini, and Dr. Liang Zhao, for their valuable comments on the manuscript. We thank the curators of CDBI, KUN, PE, and WCSBG for allowing us to use their scanned images of specimens and for research facilities. This work was supported by the National Natural Science Foundation of China (grant nos. 31870184, 31770218, and 31970210).

References

- Fei WQ, Yuan Q, Yang QE (2022) *Ranunculus huainingensis* and *R. lujiangensis* (Ranunculaceae), described from Anhui in China, are both synonymous with *R. ternatus*, a polymorphic eastern Asian species. *Phytotaxa* 573(1): 15–38. <https://doi.org/10.11646/phytotaxa.573.1.2>
- Fei WQ, Yuan Q, Yang QE (2023a) *Ranunculus jiguanishanicus* (Ranunculaceae), a new species from Sichuan, China. *PhytoKeys* 219: 57–75. <https://doi.org/10.3897/phytokeys.219.96266>
- Fei WQ, Yuan Q, Yang QE (2023b) *Ranunculus luanchuanensis* (Ranunculaceae), a new species from Henan, China. *PhytoKeys*. (in press)

- Hörandl E, Emadzade K (2012) Evolutionary classification: A case study on the diverse plant genus *Ranunculus* L. (Ranunculaceae). *Perspectives in Plant Ecology, Evolution and Systematics* 14(2): 310–324. <https://doi.org/10.1016/j.ppees.2012.04.001>
- Hörandl E, Paun O, Johansson JT, Lehnebach C, Armstrong T, Chen L, Lockhart P (2005) Phylogenetic relationships and evolutionary traits in *Ranunculus* s.l. (Ranunculaceae) inferred from ITS sequence analysis. *Molecular Phylogenetics and Evolution* 36(2): 305–327. <https://doi.org/10.1016/j.ympev.2005.02.009>
- IUCN Standards and Petitions Committee (2022) Guidelines for Using the IUCN Red List Categories and Criteria. Version 14. IUCN Standards and Petitions Committee. <https://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Levan A, Fredga K, Sandberg AA (1964) Nomenclature for centromeric position on chromosomes. *Hereditas* 52(2): 201–220. <https://doi.org/10.1111/j.1601-5223.1964.tb01953.x>
- Luo MR, Zhao L (2013) A new *Ranunculus* species (Ranunculaceae) from Shaanxi, China. *Bangladesh Journal of Plant Taxonomy* 20(2): 201–205. <https://doi.org/10.3329/bjpt.v20i2.17394>
- Paun O, Lehnebach C, Johansson JT, Lockhart P, Hörandl E (2005) Phylogenetic relationships and biogeography of *Ranunculus* and allied genera (Ranunculaceae) in the Mediterranean region and in the European Alpine System. *Taxon* 54(4): 911–930. <https://doi.org/10.2307/25065478>
- Sun H, Zhang JW, Deng T, Boufford DE (2017) Origins and evolution of plant diversity in the Hengduan Mountains, China. *Plant Diversity* 39(4): 161–166. <https://doi.org/10.1016/j.pld.2017.09.004>
- Tamura M (1995) Angiospermae. Ordnung Ranunculales. Fam. Ranunculaceae. II. Systematic Part. In: Hiepko P (Ed.) *Die Natürlichen Pflanzenfamilien* (2nd edn. Vol. 17aIV). Duncker & Humblot, Berlin, 223–519.
- Thiers B (2022) Index Herbariorum: A Global Directory of Public Herbaria and Associated Staff. <http://sweetgum.nybg.org/science/ih/> [accessed 2 December 2022]
- Wang WT (1993) *Ranunculus* L. In: Wang WT (Ed.) *Vascular Plants of the Hengduan Mountains* (Vol. 1). Science Press, Beijing, 532–541.
- Wang WT (1995a) A revision of the genus *Ranunculus* in China (I). *Bulletin of Botanical Research* 15(2): 137–180.
- Wang WT (1995b) A revision of the genus *Ranunculus* in China (II). *Bulletin of Botanical Research* 15(3): 275–329.
- Wang WT (1996) Notulae de Ranunculaceis Sinensibus (XIX). *Bulletin of Botanical Research* 16(2): 155–166.
- Wang WT (2007) *Ranunculus ailaoshanicus* W. T. Wang, a new species of Ranunculaceae from Yunnan, China. *Acta Phytotaxonomica Sinica* 45(3): 293–295. <https://doi.org/10.1360/aps06192>
- Wang WT (2008) New taxa of Ranunculaceae from Yunnan. *Acta Botanica Yunnanica*.
- Wang WT (2013) Six new species and two new varieties of Ranunculaceae from southwest China. *Guihaia* 33(5): 579–587.
- Wang WT (2015) Five new species of *Ranunculus* from west China. *Bulletin of Botanical Research* 35(5): 641–646.

- Wang WT (2016) Six new species of Ranunculaceae from China. *Guihaia* 36(11): 1303–1311.
- Wang WT (2018) *Ranunculus lujiangensis*, a new species of Ranunculaceae from Anhui Province. *Bulletin of Botanical Research* 38(6): 801–803. <https://doi.org/10.7525/j.issn.1673-5102.2018.06.001>
- Wang WT (2019a) *Ranunculus kangmaensis* W. T. Wang, a new species of Ranunculaceae from Tibet of China. *Guihaia* 39(3): 285–287. <https://doi.org/10.11931/guihaia.gxzw201712013>
- Wang WT (2019b) Two new species of *Ranunculus* from Tibet, China. *Guihaia* 39(9): 1139–1342. <https://doi.org/10.11931/guihaia.gxzw201811047>
- Wang WT (2022) Five new species and one new variety of *Ranunculus* (Ranunculaceae) from Sichuan, with one new section represented by one of these species. *Guihaia* 42(1): 1–9. <https://doi.org/10.11931/guihaia.gxzw202011047>
- Wang WT, Chen SR (2015) *Ranunculus laohegouensis*, a new species of Ranunculaceae from Sichuan. *Bulletin of Botanical Research* 38(6): 801–802. <https://doi.org/10.7525/j.issn.1673-5102.2015.06.001>
- Wang WT, Gilbert MG (2001) *Ranunculus* Linnaeus. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China* (Vol. 6). Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 391–431.
- Wang WT, Liao L (2009) *Ranunculus napahaiensis*, a new species of Ranunculaceae from Yunnan, China. *Guihaia* 29(4): 427–429.
- Wang WT, Yang Z, Xie J (2016) *Ranunculus huainingensis*, a new species of Ranunculaceae from Anhui. *Guihaia* 36(supplement 1): 97–99.
- Yang QE (2000) *Ranunculus wangianus* Q. E. Yang, a new species from NW Yunnan, China and its karyotype. *Acta Phytotaxonomica Sinica* 38(6): 551–556.
- Yuan Q, Yang QE (2017a) *Ranunculus angustisepalus* (Ranunculaceae) is an *Oxygraphis* and conspecific with *O. delavayi*. *Phytotaxa* 319(1): 103–110. <https://doi.org/10.11646/phytotaxa.319.1.6>
- Yuan Q, Yang QE (2017b) The identity of *Ranunculus ailaoshanicus* (Ranunculaceae) from China. *Phytotaxa* 319(1): 111–117. <https://doi.org/10.11646/phytotaxa.319.1.7>
- Yuan Q, Yang QE (2017c) The identity of *Ranunculus laohegouensis* (Ranunculaceae) from Sichuan, China. *Phytotaxa* 324(2): 198–200. <https://doi.org/10.11646/phytotaxa.324.2.9>
- Zhang DC, Zhang YH, Boufford DE, Sun H (2009) Elevational patterns of species richness and endemism for some important taxa in the Hengduan Mountains, southwestern China. *Biodiversity and Conservation* 18(3): 699–716. <https://doi.org/10.1007/s10531-008-9534-x>
- Zhang M, Zeng YP, Yang QE (2020) *Ranunculus kangmaensis* (Ranunculaceae), a new synonym of *Halerpestes tricuspis* var. *variifolia*. *Phytotaxa* 434(1): 101–112. <https://doi.org/10.11646/phytotaxa.434.1.7>